

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

P402 Interstitial and granulomatous lung disease in inflammatory bowel disease patients

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1659679> since 2020-07-11T10:37:34Z

Publisher:

Oxford University Press

Published version:

DOI:10.1093/ecco-jcc/jjx180.529

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

Conclusions: Our results suggest that appropriate dose adjustments by reducing the intervals of IFX dosing may overcome low ADA titers with the aim to maintain or restore clinical response. In addition, proactive evaluation of ADA status may allow earlier identification of patients prone to a loss of response. The implementation of Bayesian dashboards may assist in the prediction of SICs <3 µg and determining adequate dosing modifications.

P402

Interstitial and granulomatous lung disease in inflammatory bowel disease patients

E. Eliadou^{1*}, J. Moleiro², D.G. Ribaldone³, M. Astegiano³, K. Rothfuß⁴, C. Taxonera⁵, B. Verstockt⁶, F. Ghalim⁷, F. Carbonnel⁷, S. Festa⁸, A. Berrozpe⁹, L. Maia¹⁰, E. Savarino¹¹, P. Ellul¹², E. Zagorowicz¹³, S. Vavricka¹⁴, M. Calvo¹⁵, I. Koutroubakis¹⁶, F. Hoentjen¹⁷, L. Fernández Salazar¹⁸, F. Callela¹⁹, F. Cañete Pizarro²⁰, K. Soufleris²¹, E. Sonnenberg²², M. Cavicchi²³, G. Fiorino^{24,25}

¹Manchester Royal Infirmary, Department of Gastroenterology, Manchester, UK, ²Instituto Portugues de Oncologia de Lisboa Francisco Gentil, Lisboa, Portugal, ³Gastroenterologia-U, Città della Salute e della Scienza di Torino, Turin, Italy, ⁴Robert-Bosch Hospital, Gastroenterology Department, Stuttgart, Germany, ⁵Servicio Aparato Digestivo, Hospital Clínico San Carlos, IBD unit, Madrid, Spain, ⁶University Hospitals Leuven, Gastroenterology Department, Leuven, Belgium, ⁷Kremlin Bicêtre Hospital, AP-HP, University Paris Sud, Gastroenterology Department, Paris, France, ⁸Ospedale S. Filippo Neri, Gastroenterology Department, Rome, Italy, ⁹Bellvitge's hospital, Gastroenterology Department, Barcelona, Spain, ¹⁰Centro Hospitalar do Porto, Gastroenterology Department, Porto, Portugal, ¹¹University of Padua, Padua, Italy, ¹²Mater Dei Hospital, Msida, Malta, ¹³The Maria Skłodowska Curie Memorial Cancer Centre and Institute of Oncology, Gastroenterology Department, Warsaw, Poland, ¹⁴Triemli Hospital, Zurich, Switzerland, ¹⁵Hospital Universitario Puerta de Hierro Madrid, Madrid, Spain, ¹⁶University of Crete, Heraklion, Greece, ¹⁷Radboud University Medical Center, Nijmegen, The Netherlands, ¹⁸Clínico Universitario de Valladolid, Valladolid, Spain, ¹⁹UOC Gastroenterologia, Ospedale San Giuseppe, Florence, Italy, ²⁰Hospital Universitari Germans Trias i Pujol in Badalona, Barcelona, Spain, ²¹Theagenion Cancer Hospital of Thessaloniki, Thessaloniki, Greece, ²²Charité-Universitätsmedizin Berlin, Berlin, Germany, ²³Clinique de Bercy, Paris, France, ²⁴RCCS Istituto Clinico Humanitas, Gastroenterology Department, Rozzano, Milan, Italy, ²⁵For ECCO CONFER Steering Committee, Vienna, Austria

Background: Granulomatous (GL) and interstitial lung disease (ILD) are rare respiratory disorders that have been associated with inflammatory bowel disease (IBD). Clinical presentation is polymorphic and aetiology is unclear.

Methods: This was European Crohn's and colitis organisation (ECCO) retrospective observational study performed as part of CONFER project. A call to all ECCO members was made to report concomitant granulomatous or ILD and IBD cases. Clinical data were recorded in a standardised case report form.

Results: Twenty-two granulomatous lung disease patients were identified from 18 university hospitals, 17 males and 5 females with a mean age of 46 years (18–86); 17 patients with Crohn's disease (CD) and five with ulcerative colitis (UC). In 19 patients IBD diagnosis preceded lung disease in a median time of 10.6 years (0–27). In three patients lung disease were diagnosed in a median time of 14 months

(8–24) before IBD. Only four patients had active disease. Seven patients had drug-related granulomatous disease (sarcoidosis $n = 4$) and 14 had non-drug-related GL (primary sarcoidosis $n = 7$, fungal infection $n = 2$ and unspecified $n = 5$). Ten patients (45%) required hospitalisation but none required invasive ventilation. Fifteen of 22 patients received systemic steroids and causative drug was stopped in all patients. At further follow-up, 15 of 22 patients had no respiratory symptoms. Thirty-one patients with ILD were identified from 14 medical centres, 12 females and 19 males with mean age of 47 years (17–84); eight patients had CD, 22 had UC and one had indeterminate colitis. All patients had IBD diagnosis prior to ILD with a median time of 10.27 years (0.3–51). Eight patients had active disease. Eleven patients had non-drug-related ILD and 20 had drug-related ILD (mesalazine $n = 9$, methotrexate $n = 1$, golimumab $n = 1$, vedolizumab $n = 1$ and infliximab $n = 8$). ILD cases were classified as: Cryptogenic organising pneumonia $n = 11$, Eosinophilic pneumonia $n = 2$, bronchiolitis $n = 2$, acute interstitial pneumonia $n = 8$, interstitial lung disease due to connective tissue $n = 1$, idiopathic fibrosis $n = 6$ and unclassified $n = 1$. Twenty-five patients (80.6%) required hospitalisation and one required non-invasive ventilation. Twenty-seven patients (87%) received systemic steroids and causative drug was stopped in all patients. At further follow-up, 14 of 31 patients had no respiratory symptoms, 7 of 31 had some improvement, and 4 of 31 had ongoing symptoms. There was one patient referred for consideration of lung transplant due to progressive fibrosis and one death due to mesalazine.

Conclusions: GL and ILD are rare, but require hospitalisation and systemic steroids. In our case series half of cases were drug-related but there was no signal of relationship between IBD therapy and the onset of ILD. More studies are needed to investigate the pathogenesis and causative association.

P403

Treatment naïve newly diagnosed patients with Crohn's disease have microbial dysbiosis correlated with disease activity and faecal calprotectin—results from a prospective inception cohort

H. Yanai^{1,2*}, I. Goren^{3,4}, L. Reshef⁵, L. Godny^{1,2}, K. Yadgar¹, K. Zonenesain¹, I. Dotan^{2,3}

¹IBD Center, Division of Gastroenterology, Rabin Medical Center, Petah Tikva, Israel, ²Sackler Faculty of Medicine, Tel Aviv Israel, Tel Aviv, Israel, ³Rabin Medical Center, Division of Gastroenterology, Petah Tikva, Israel, ⁴Sackler Faculty of Medicine, Tel Aviv University, Tel Aviv, Israel, ⁵Department of Molecular Microbiology and Biotechnology, George S. Wise Faculty of Life Sciences, Tel Aviv University, Tel Aviv, Israel

Background: Microbial dysbiosis is believed to play a role in Crohn's disease (CD). Most data are derived from CD patients under medications, an exposure that might impact microbial composition. Our aim was to assess microbial dysbiosis in patients with newly diagnosed CD and correlate it with disease activity.

Methods: Newly diagnosed CD patients were prospectively recruited and followed longitudinally. Clinical data, disease activity (physician global assessment [PGA] ranging from 0 to 3), and serum and faecal inflammatory biomarkers, were collected. Faecal samples were assessed for microbial composition using 16S rRNA sequencing. The microbial dysbiosis index (MDI) was used to quantify the degree of dysbiosis per sample.